#### THIS EVALUATION WAS FUNDED BY THE EPISCOPAL HEALTH FOUNDATION AND PROSPERA HOUSING COMMUNITY SERVICES.

#### **DECEMBER 2022**

# A QUASI-EXPERIMENTAL STUDY OF A COLLABORATIVE SERVICE MODEL BETWEEN A SUPPORTIVE HOUSING PROVIDER AND A MANAGED CARE ORGANIZATION



#### PREPARED AND PRESENTED BY

UNIVERSITY OF TEXAS HEALTH SCIENCE CENTER AT HOUSTON (UTHEALTH) SCHOOL OF PUBLIC HEALTH JACK TSAI, PHD, MSCP SUJA RAJAN, PHD CHAU TRUONG, MPH VANESSA SCHICK, PHD CECILIA GANDUGLIA CAZABAN, MD, DRPH



#### **ACKNOWLEDGEMENTS**



THANKS TO...

#### **Episcopal Health Foundation:**

Shao-Chee Sim and C.J. Eisenbarth Hager

#### Prospera Housing Community Services:

Scott Ackerson, Carmen Hancock, Diane Warren, staff and the residents

#### **Superior HealthPlan:**

Michelle Murdock, Alex Goldson, and analysts



## **EXECUTIVE SUMMARY**

Many managed care organizations (MCOs) are now focused on screening and addressing social determinants of health. Unique partnerships between MCOs with community providers are being explored but there has been little formal evaluation of these partnerships. In this study, we evaluated a collaborative, community-based service model in which participants had healthcare coverage through Superior HealthPlan, the largest MCO in Texas, and lived in properties maintained by Prospera Housing Community Services, an affordable housing provider. This Prospera+Superior collaborative model allowed for facilitated care and joint programs aimed to improve social determinants of health, including access to nutritious foods, transportation, affordable healthcare, and secure housing.

Using a quasi-experimental two-groups research design, we compared a sample of 104 participants served by the Prospera+Superior collaborative model to a demographically matched group of 104 participants who had healthcare coverage through the Superior HealthPlan Medicaid MCO but did not live at Prospera properties (i.e., Superior Only group). The primary outcomes were healthcare utilization and costs. We analyzed data from medical claims to examine change in outcomes from 12 months before implementation of the Prospera+Superior collaborative model in 2019 to 12 months after implementation. We conducted regression-based adjusted analyses which revealed the Prospera+Superior group had a 17% lower rate of emergency department/urgent care visits and spent \$2,061 less than the Superior Only group after implementation. Together, these findings provide needed evidence of the clinical and economic value of forming multi-sector collaborative models between MCOs and other community providers.

### **BACKGROUND**

There is widespread interest from the medical and healthcare field to address social determinants of health. The housing and communities in which people live can impact their health and access to healthcare. There are well-documented links between income, housing, and health. For example, financial hardship is a robust predictor of health among low-income housing residents (1) and financial strain may mediate the link between mental illness and homelessness (2). However, there is a need to examine policy levers and service models that comprehensively and effectively improve financial health, housing stability, and access to healthcare services.

In the healthcare landscape of the U.S., managed care organizations (MCOs) play an essential role in supporting the health of low-income and disabled populations. Many states are beginning to require MCOs to screen for and address social determinants of health among enrollees (3, 4). However, despite experimenting with various diverse programs to address social determinants of health, many MCOs are struggling with service integration, financing, and evaluation efforts to determine the effectiveness and sustainability of these programs (5).


Many low-income individuals face challenges with obtaining and maintaining stable housing. Supportive housing programs which provide subsidized housing are a promising housing option for individuals who need long-term support. However, as concluded in a report by the National Academies of Sciences, Engineering, and Medicine, there is no substantial evidence as yet that supportive housing programs improve health outcomes (6). In other words, supportive housing programs improve housing outcomes (7, 8), but may not necessarily improve health (6, 9).



Collaborative service models that combine healthcare and homeless services have shown promise. For example, federally-funded Health Care for the Homeless (HCH) clinics around the country serve as primary care "health homes" for individuals experiencing and at-risk of homelessness and have grown to over 200 sites over the past 3 decades (10). The U.S. Department of Veterans Affairs (VA) has implemented the Homeless Patient Aligned Care Team (H-PACT) program which is a multidisciplinary medical care home that offers tailored primary care services to homeless veterans. There is evidence that use of H-PACT is associated with reductions in emergency department (ED) use, improvements in primary care utilization, and positive patient experiences (11). However, most of the existing models have largely been hospitalbased instead of community-based.

Unique collaborative models can offer clients multiple services on-site and provide a variety of partnered services. These models can also increase communication between different providers and entities which can enhance care and improve continuity of care. Experts have argued for over a decade for innovative ways to integrate health insurance and housing services. A well-known policy brief that was disseminated over a decade ago made a business case for Medicaid-financed services in supportive housing to lower costs associated with avoidable hospitalizations and other crisis services (12). There is a need to empirically study new, innovative community-based service models between housing providers and insurance payers.

#### PROJECT AIMS



In the current study, we conducted a quasi-experimental study to examine a collaborative, community-based service model that involved a supportive housing provider and a MCO compared to a comparison group enrolled in the MCO. Our primary outcomes were use of healthcare services and costs.

WE HYPOTHESIZED
THAT THE
COLLABORATIVE
SERVICE MODEL
WOULD BE
ASSOCIATED WITH
LESS USE OF
ACUTE CARE AND
LOWER OVERALL
MEDICAL CARE
COSTS THAN A
COMPARISON
GROUP.

## **METHODS**







RESEARCH DESIGN DATA SOURCES






MEASURES



**DATA ANALYSIS** 




#### PROGRAM DESCRIPTION

Prospera Housing Community Services is an organization that has been building and operating supportive housing in Texas for over 25 years. Permanent supportive housing models provide long-term housing and supportive services to individuals without time limits or retributions regardless of service engagement. Prospera Housing Community Services serves individuals and families by providing safe, highquality, affordable housing and support services at over 50 sites across 19 cities throughout South and Central Texas. At each site, there are on-site service managers that help offer resident services and facilitate care and billing of services with insurance payers like Superior HealthPlan. In this study, residents across 11 sites were included and these sites are listed in Appendix A.

Superior HealthPlan was founded in 1999 in El Paso, Texas and has become the largest MCO in Texas. Superior HealthPlan operates under the parent company Centene. Superior HealthPlan provides enrollees with access to Medicaid STAR, STAR+PLUS, STAR Health, STAR Kids and CHIP, Medicare Advantage, and the Health Insurance Marketplace.

A collaborative service model involving the affordable housing provider, Prospera Housing Community Services, and the Superior HealthPlan Medicaid MCO, was implemented in 2019. This partnership allowed for facilitated care between the two organizations and joint programs that aimed to improve social determinants of health, including access to nutritious foods, transportation, affordable healthcare, and secure housing (13). Over 50 programs were offered through this partnership, which are detailed in Appendix B. This study focused on testing this partnership holistically as a collaborative service model and not individual programs.




#### RESEARCH DESIGN

This study used a matched-groups design (matching on age, gender, service delivery area, and have 6 months of coverage) combined with regression-based adjusted analyses to compare a group of 104 residents living at properties at Prospera Housing Community Services with coverage by Superior HealthPlan (herein referred to at Prospera+Superior) to a matched group of 104 residents who did not live at Prospera Housing Community Services and only had coverage by Superior HealthPlan (herein referred to as Superior Only). Prior to the regression analysis, participants in the Prospera+Superior group were matched with similar individuals from the Superior group, to develop a comparable control group. These groups were matched on demographics (age, gender service delivery area), any dual coverage (e.g., Medicaid-Medicare coverage), and member product (e.g., Star+Plus).

To test our study hypothesis, we examined the Prospera+Superior and Superior Only groups in the 12 months before implementation of the Prospera+Superior collaboration (i.e., before 2019) and the 12 months after implementation. The main outcomes were healthcare utilization and costs.

All study procedures were approved by the institutional review board at the University of Texas Health Science Center at Houston (Project # HSC-SPH-21-0841).




#### DATA SOURCES

Data from enrollment, medical and pharmacy files in 2018-2020 from Superior HealthPlan for both Prospera+Superior and Superior Only groups were transferred to the research team at UTHealth Center for Health Care Data. Among dual eligible participants, some were those enrolled in the Medicaid-Medicare Managed Care waiver program (MMP) and some were not. Full claims history data were received for those in the MMP program; for dual eligible participants not in the MMP, data were pulled from Medicare claims files available through the UTHealth Center for Health Care Data, which is a Centers for Medicare & Medicaid Services (CMS) Qualified Entity. This linkage allowed for a more complete review of healthcare utilization and costs for the study samples.

To ensure there was no overcounting of services and costs, manual review of data from randomly selected participants was conducted and compared across different data sources. The analytic dataset created for this study contained information on participants' registration in Prospera housing (for the Prospera+Superior group), demographic characteristics, Medicaid enrollment information, healthcare utilization (i.e., emergency department visits, inpatient admissions, outpatient visits) and spending for healthcare services.

The first day the Prospera+Superior collaboration was implemented at each housing site in 2019 served as the index date and the analytic dataset included the 12 months before and after the index date.



#### **MEASURES**

Information on demographic characteristics and healthcare coverage of participants were extracted from eligibility and enrollment files.

Medical claims were reviewed to examine medical diagnoses and to measure each participant's Charlson Comorbidity Index (CCI), a method of categorizing and weighting comorbidities to predict mortality risk (14).

Four healthcare utilization measures and two types of healthcare costs were examined before and after implementation of the Prospera+Superior collaboration to study the effect of the collaborative service model on participants. These measures were, outpatient visits, ED/urgent care visits, inpatient visits and inpatient length of stay, medical care costs, and pharmaceutical costs.



First, the Prospera+Superior and Superior Only groups were compared on demographics, healthcare coverage, and clinical diagnoses using bivariate tests with independent t-tests and chisquare tests.

Second, the groups were compared descriptively on healthcare utilization and costs before and after implementation of the Prospera+Superior collaboration. Wilcoxon signed rank test was used to compare the average mean between groups unadjusted for pre-implementation differences.

Third, the groups were compared on healthcare utilization and costs controlling for differences in healthcare coverage and clinical diagnoses before implementation of the Prospera+Superior collaboration.


Since the data has a panel structure where the patients are repeat sampled before and after implementation, panel data regressions were used based on various specification tests to help pick the right regressions. The fixed effect panel data regression analysis was used, which is the most conservative regression method in this context. For healthcare utilizations, which had large ranges, a fixed effects linear regression was used. For healthcare utilizations, such as ED/urgent care visits and number of inpatient admissions, which had a limited range, fixed effects Poisson regression was used to account for the "count" data-like nature of the utilization measures. We tried other mixed effects models, and our results were robust to the type of regression used.

#### RESULTS - DEMOGRAPHIC DIFFERENCES





There were no significant demographic differences between the Prospera+Superior and Superior groups.




#### RESULTS - MEMBER PRODUCT



#### **Superior Only**


THOSE IN THE PROSPERA +
SUPERIOR GROUP WERE
SIGNIFICANTLY MORE
LIKELY TO HAVE STAR+
THAN THOSE IN THE
SUPERIOR ONLY GROUP.




**Prospera + Superior** 

#### CHARLSON COMORBIDITY INDEX (CCI)

The most common medical conditions were diabetes with and without complications, and renal disease.



#### **Mean # of CCI Conditions**



The Superior Only group had more CCI conditions than the Prospera+ Superior group.

#### HEALTHCARE UTILIZATION

Healthcare utilization before (pre) and after (post) implementation of the collaborative model intervention in the Prospera+Superior group.

Healthcare Visits	Pre- implementation Difference	Post- implementation Difference
# inpatient admissions	-	0
Inpatient length of stay	-	0
# of ED/urgent care vists	-	-
# of outpatient visits	0	0

<sup>-</sup> No difference between groups at timepoint.

Utilization lower in Prospera + Superior group than Superior Only group.

After implementation, the Prospera + Superior Only group <u>continued</u> to have significantly lower outpatient & inpatient utilization.



THE PROSPERA +
SUPERIOR
GROUP HAD A
17% LOWER
RATE OF
ED/URGENT
CARE VISITS
RELATIVE TO
THE SUPERIOR
ONLY GROUP.

## PHARMACEUTICAL COSTS

Pharmaceutical costs before (pre) and after (post) implementation of the collaborative model intervention in the Prospera+Superior group.

Costs	Pre- implementation Difference	Post- implementation Difference
Medical Cost Dollars	-	-
Pharmacy Cost Dollars	-	0

- No difference between groups at timepoint.
- Costs lower in Prospera + Superior group than Superior Only group.

The Prospera+Superior group spent

\$2,061

less than the Superior Only group after implementation of the Prospera+Superior collaborative model after controlling for preimplementation differences.

#### DISCUSSION

There has been wide interest in multi-sector collaborative models of care, particularly between MCO and community providers. However, there has been limited empirical data to support the effectiveness of particular models. This study contributes to the literature by controlled comparison between a collaborative service model partnering an MCO and an permanent supportive housing provider. Our main finding showed this model was associated with decreased use of ED/urgent care services compared to a group that was under the same Medicaid MCO but did not have the opportunities of the collaborative services with a housing provider. While participants were not randomized so we cannot infer causality, the finding does suggest the collaborative model improved access to care and facilitated greater participation in healthcare prevention activities upstream that resulted in fewer acute care needs.

#### **FINDINGS**

A secondary and important finding was that the collaborative service model was associated with lower overall pharmaceutical costs among its participants than a comparison group. This finding provides further data in supporting a business case for these type of collaborative models (12). Furthermore, given the wide reliance on medications and efforts to reduce inappropriate polypharmacy (15, 16) as well as concerns about rising medication costs in the U.S. (17), the observed decrease in pharmaceutical costs may have broad program and policy implications. For example, one study of the top 150 medications administered and prescribed in EDs in the U.S. found the costs increased by 28-125% over the past decade. It may be important to note that while our findings support other collaborative efforts to integrate care in different settings, the Prospera+Superior collaborative model is different from more medicallybased models such as the National Health Care for the Homeless clinics (10) and the VA H-PACT programs (11). The Prospera+Superior model is embedded in where people live and works to address multiple social determinants of health instead of specific health conditions.




#### LIMITATIONS



There were several limitations of the study worth noting. First, as mentioned earlier, we did not randomize participants into groups, so a full randomized clinical trial is needed to confirm these findings. Second, this was a unique partnership between Prospera and Superior HealthPlan and the generalizability of these findings to partnerships with other similar agencies has yet to be determined. Third, given our sample size, we were limited to examining broad categories of healthcare utilization and costs and did not have adequate cell sizes to examine more specific categories as originally intended (e.g., use of specialized preventive services or costs). Further multi-site studies are needed to examine these issues more directly.

THESE LIMITATIONS WERE **COUNTERBALANCED BY** THE STRENGTHS OF THE STUDY. WHICH **INCLUDED A RESEARCH DESIGN WITH A COMPARISON GROUP, A** STATISTICALLY **RIGOROUS APPROACH** THAT INCLUDED **MATCHED GROUPS WITH REGRESSION-BASED ADJUSTED ANALYSES.** AND EXAMINATION OF **BOTH HEALTHCARE UTILIZATION AND** COSTS.

#### CONCLUSIONS



## Service Models

There is evidence that collaborative service models between MCOs and housing providers, like between Superior HealthPlan and Prospera Housing Community Services, can reduce use of costly ED and urgent care services and overall pharmaceutical costs. These findings may have policy implications as MCOs focus on social determinants of health and consider new models of care to effectively address them.



#### REFERENCES

- 1. Tucker-Seeley RD, Harley AE, Stoddard AM, Sorensen GG. Financial hardship and self-rated health among low-income housing residents. Health Education & Behavior. 2013;40(4):442-8.
- 2. Elbogen EB, Lanier M, Wagner HR, Tsai J. Financial strain, mental illness, and homelessness: Results from a national longitudinal study. Medical Care. 2021;59:S132-S8.
- 3. Shrank WH, Keyser DJ, Lovelace JG. Redistributing investment in health and social services—the evolving role of managed care. JAMA. 2018;320(21):2197-8.
- 4. Health Management Associates. Medicaid managed care provides opportunities for states to address social determinants of health and health equity. Health Management Associates; 2022 [cited 2022 October 27]; Available from: https://www.healthmanagement.com/blog/medicaid-managed-care-provides-opportunities-for-states-to-address-social-determinants-of-health-and-health-equity/.
- 5. Gottlieb L, Ackerman S, Wing H, Manchanda R. Understanding Medicaid managed care investments in members' social determinants of health. Population Health Management. 2017;20(4):302-8.
- 6. National Academies of Sciences E, and Medicine. Permanent supportive housing: Evaluating the evidence for improving health outcomes among people experiencing chronic homelessness. Washington, DC: The National Academies Press2018.
- 7. Tsai J, Mares AS, Rosenheck RA. A multi-site comparison of supported housing for chronically homeless adults: "Housing first" versus "residential treatment first". Psychological Services. 2010;7(4):219-32.
- 8. Leff HS, Chow CM, Pepin R, Conley J, Allen IE, Seaman CA. Does one size fit all? What we can and can't learn from a meta-analysis of housing models for persons with mental illness. Psychiatric Services. 2009;60:473-82.
- 9. Tsai J, Gelberg L, Rosenheck RA. Changes in physical health after supported housing: Results from the collaborative initiative to end chronic homelessness. Journal of General Internal Medicine. 2019:34(9):1703-8.
- 10. Zlotnick C, Zerger S, Wolfe PB. Health care for the homeless: What we have learned in the past 30 years and what's next. American Journal of Public Health. 2013;103(Suppl 2):S199-S205.
- 11. Tsai J, Havlik J, Howell BA, Johnson E, Rosenthal D. Primary care for veterans experiencing homelessness: A narrative review of the Homeless Patient Aligned Care Team (HPACT) model. Journal of General Internal Medicine. 2022.
- 12. Nardone M, Cho R, Moses K. Medicaid-financed services in supportive housing for high-need homeless beneficiaries: The business case. Hamilton, NJ: Center for Health Care Strategies, Inc.2012.
- 13. Services PCC. Superior HealthPlan. San Antonio, TX: Prospera Housing and Community Services; 2015 [cited 2022 December 4]; Available from: https://prosperahcs.org/partners/superior-healthplan/.
- 14. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. Journal of Chronic Diseases. 1987;40(5):373-83.
- 15. Scott IA, Hilmer SN, Reeve E, Potter K, Le Couteur D, Rigby D, et al. Reducing inappropriate polypharmacy: The process of deprescribing. JAMA Internal medicine. 2015;175(5):827-34.
- 16. Reeve E, Shakib S, Hendrix I, Roberts MS, Wiese MD. Review of deprescribing processes and development of an evidence-based, patient-centred deprescribing process. British Journal of Clinical Pharmacology. 2014;78(4):738-47.
- 17. Leighl NB, Nirmalakumar S, Ezeife DA, Gyawali B. An arm and a leg: the rising cost of cancer drugs and impact on access. American Society of Clinical Oncology Educational Book. 2021;41:e1-e12.

## APPENDIX A

#### LIST OF PROSPERA PROPERTIES IN THE STUDY

Property Name	# of residents in the study	Start date of Prospera + Superior partnership		
	n= 104	partnership		
Country Club	5 (5%)	1/1/2019		
Hacienda Senior	10 (10%)	1/1/2019		
Kingsville LULAC	4 (4%)	1/1/2019		
La Risa	15 (14%)	1/1/2019		
Laredo Manor	7 (7%)	4/30/2019		
Las Palmas Gardens	8 (8%)	12/1/2019		
Oak Manor	17 (16%)	4/30/2019		
Palms at Leopard	10 (10%)	12/1/2019		
Terraces at Haven	5 (5%)	12/1/2019		
Vista Verde	17 (16%)	4/30/2019		
Woodland Creek	6 (6%)	4/30/2019		

## APPENDIX B

PROGRAMS OFFERED THROUGH PROSPERA-SUPERIOR HEALTHPLAN PARTNERSHIP, SORTED BY SOCIAL DETERMINANTS OF HEALTH

#### APPENDIX C- TABLE 1.

Table 1. Demographic and coverage characteristics of Prospera+Superior and Superior Only groups

Total Members	Superior O n= 1		Prospera+Superior group n=104		<i>p</i> -value	
	Mean/Count	SD/%	Mean/Count	SD/%		
Age	59.96	17.99	57.22	15.46	0.240	
Gender						
Female	60	58%	60	58%		
Male	44	42%	44	42%		
<b>Dual Status</b>						
No	42	40%	50	48%	0.064	
Yes	62	60%	54	52%	0.264	
Member Product						
MMP	44	42%	2	2%	<0.001	
Star+Plus	60	58%	102	98%	< 0.001	
Race/Ethnicity						
Asian/Pacific Islander	2	2%	1	1%		
Black (Non-Hispanic)	6	6%	9	9%		
White (Non-Hispanic)	16	15%	10	10%	0.301	
Hispanic	57	55%	69	66%	0.301	
Other	1	1%	0	0%		
Not Provided	22	21%	15	14%		
Prospera properties						
Country Club	N/A	N/A	5	4.8%	N/A	
Hacienda Senior	N/A	N/A	10	9.62%	N/A	
Kingsville LULAC	N/A	N/A	4	3.85%	N/A	
La Risa	N/A	N/A	15	14.42%	N/A	
Laredo Manor	N/A	N/A	7	6.73%	N/A	
Las Palmas Gardens	N/A	N/A	8	7.69%	N/A	
Oak Manor	N/A	N/A	17	16.35%	N/A	
Palms at Leopard	N/A	N/A	10	9.62%	N/A	
Terraces at Haven	N/A	N/A	5	4.81%	N/A	
Vista Verde	N/A	N/A	17	16.35%	N/A	
Woodland Creek	N/A	N/A	6	5.77%	N/A	

Note: Prospera+Superior group lived on Prospera properties and had coverage through Superior HealthPlan while the Superior Only group did not live on Prospera properties but had Superior HealthPlan coverage. MMP= Medicare-Medicaid Plan. Star+Plus= Texas Medicaid managed care program. N/A= Not applicable. Pink boxes indicate statistically significant values.

Table 1 shows the demographic characteristics and healthcare coverage of participants. The majority of participants were female, Hispanic, and had dual healthcare coverage. There were no significant differences between the Prospera+Superior and Superior Only groups on demographic characteristics and rates of dual coverage, which was expected given the two groups were matched on these characteristics. However, the Prospera+Superior group was more likely to be in the Star+Plus program and less likely to be in the MMP program than the Superior Only group.

#### APPENDIX D- TABLE 2. MEDICAL CONDITIONS

Table 2 shows the CCI's medical conditions among participants. The most common medical conditions were diabetes with and without complications, and renal disease.

Table 2. Medical conditions from the Charlson Comorbiditiy Index (CCI) among Prospera+Superior and Superior Only groups

<u>CCI</u>	Superior Only		Prospera+Superior		<i>p</i> -value
	<u>N</u>	%	N	%	
Congestive Heart Failure	11	11%	2	2%	0.010
Peripheral Vascular Disease	17	16%	3	3%	0.001
Cerebrovascular Disease	12	12%	4	4%	0.037
Diabetes w/o complications	42	40%	22	21%	0.003
Diabetes w complications	37	36%	16	15%	0.001
Renal Disease	27	26%	8	8%	< 0.001
Mean Average CCI Score (SD)	4.40 (3.63)		2.89 (2.45)		0.001

Note: Prospera+Superior group lived on Prospera properties and had coverage through Superior HealthPlan while the Superior Only group did not live on Prospera properties but had Superior HealthPlan coverage. Pink boxes indicate statistically significant values.

The Superior Only group was significantly more likely to have six medical conditions from the CCI and had higher CCI scores than the Prospera+Superior group.

#### APPENDIX E- TABLE 3. UTILIZATION

Table 3 shows the healthcare utilization of inpatient, outpatient, and ED/urgent care services 12 months before and after implementation of the Prospera+Superior collaborative model among the Prospera+Superior and Superior Only groups.

Table 3. Unadjusted healthcare utilization and cost measures between the Prospera+Superior and Superior Only groups pre and postimplementation of the Prospera+Superior collaboration

	Pre-implementation period			Post-implementation period				
Utilization Measure (Mean, SD)	Superior Only	Prospera+ Superior	Diff	p-value	Superior Only	Prospera+ Superior	Diff	p-value
# of inpatient admissions	0.28 (1.07)	0.09 (0.46)	+0.19	0.071	0.27 (0.87)	0.05 (0.32)	+0.22	0.008
Inpatient length of stay	3.81 (19.09)	0.58 (3.79)	+3.23	0.052	1.96 (8.48)	0.13 (0.86)	+1.83	0.003
# of ED/urgent care visits	0.42 (0.94)	0.50 (1.21)	-0.08	0.948	0.56 (1.34)	0.29 (0.77)	+0.27	0.081
# of outpatient visits	6.49 (14.02)	4.66 (11.52)	+1.83	0.014	7.85 (18.55)	4.54 (11.82)	+3.31	0.020
Medical cost dollars	6,508 (17,007)	6,190 (13,934)	+318	0.947	7,398 (14101)	6,297 (10,469)	+1,101	0.509
Pharmacy cost dollars	1,214 (2,913)	3,103 (9,053)	-1,889	0.109	2,656 (5210)	2,483 (8,696)	+173	0.041

Note: Prospera+Superior group lived on Prospera properties and had coverage through Superior HealthPlan while the Superior Only group did not live on Prospera properties but had Superior HealthPlan coverage. ED= Emergency Department. Pink boxes indicate statistically significant values.

Before implementation, the Superior Only group had significantly higher outpatient utilization than the Prospera+Superior group, which is consistent with the higher number of medical conditions found in the Superior Only group. After implementation, the Superior Only group continued to have significantly higher outpatient as well as inpatient utilization and higher pharmaceutical costs than the Prospera+Superior group.

#### APPENDIX F- TABLE 4. MULTIVARIABLE ANALYSES

Due to group differences in background and clinical characteristics before implementation, multivariable analyses were conducted controlling for these differences to examine differences between groups on healthcare utilization.

Table 4. Multivariable comparisons of Prospera+Superior and Superior only groups on healthcare utilization using fixed effects panel data regression analyses

Type of Service Utilization	Effect Size	95% Confidence Interval
Outpatient visits	-1.48	-4.88, 1.92
Emergency department/urgent care visits	-0.83†	-1.42, -0.23
Inpatient admissions	-0.43†	-1.67, 0.80
Length of stay for inpatient admissions	1.40	-2.10, 4.90
Medical care costs	-782.61	-5,107.51, 3,542.29
Prescription medication costs	-2,061.24	-3,219.22, -903.26

Note: Prospera+Superior group lived on Prospera properties and had coverage through Superior HealthPlan while the Superior Only group did not live on Prospera properties but had Superior HealthPlan coverage. Effect sizes are marginal effects based on fixed effects linear regressions, except effect sizes denoted with † are incidence rate ratios based on fixed effects Poisson regression used for count data. Pink boxes indicate statistically significant values at p<.05 level.

As shown in Table 4, the Prospera+Superior group had a significant 17% lower rate of ED/urgent care visits than the Superior Only group after controlling for differences in baseline utilization and clinical characteristics. The Prospera+Superior group also had significantly lower pharmaceutical costs than the Superior Only group.

THIS EVALUATION WAS FUNDED BY THE EPISCOPAL HEALTH FOUNDATION AND PROSPERA HOUSING COMMUNITY SERVICES.

#### **SUGGESTED CITATION**

TSAI J, RAJAN S, TRUONG C, SCHICK V, & GANDUGLIA C. (2022). A QUASI-EXPERIMENTAL STUDY OF A COLLABORATIVE SERVICE MODEL BETWEEN A SUPPORTIVE HOUSING PROVIDER AND A MANAGED CARE ORGANIZATION. HOUSTON, TX. A PRELIMINARY REPORT FOR THE EPISCOPAL HEALTH FOUNDATION.

